

EQUIVARIANT COHOMOLOGY AND ANALYTIC DESCRIPTIONS OF RING ISOMORPHISMS

BO CHEN AND ZHI LÜ

ABSTRACT. In this paper we consider a class of connected closed G -manifolds with a non-empty finite fixed point set, each M of which is totally non-homologous to zero in M_G (or G -equivariantly formal), where $G = \mathbb{Z}_2$. With the help of the equivariant index, we give an explicit description of the equivariant cohomology of such a G -manifold in terms of algebra, so that we can obtain analytic descriptions of ring isomorphisms among equivariant cohomology rings of such G -manifolds, and a necessary and sufficient condition that the equivariant cohomology rings of such two G -manifolds are isomorphic. This also leads us to analyze how many there are equivariant cohomology rings up to isomorphism for such G -manifolds in 2- and 3-dimensional cases.

1. INTRODUCTION

Throughout this paper, assume that $G = \mathbb{Z}_2$ unless stated otherwise. Let $EG \rightarrow BG$ be the universal principal G -bundle, where $BG = EG/G = \mathbb{R}P^\infty$ is the classifying space of G . It is well-known that $H^*(BG; \mathbb{Z}_2) = \mathbb{Z}_2[t]$ with the t one-dimensional generator.

Let X be a G -space. Then $X_G := EG \times_G X$ —the orbit space of the diagonal action on the product $EG \times X$ —is the total space of the bundle $X \rightarrow X_G \rightarrow BG$ associated to the universal principal bundle $G \rightarrow EG \rightarrow BG$. The space $X_G = EG \times_G X$ is called the *Borel construction* on the G -space X . Applying cohomology with coefficients $\mathbb{Z}_2$ to X_G gives the *equivariant cohomology* $H_G^*(X; \mathbb{Z}_2) := H^*(X_G; \mathbb{Z}_2)$. It is well-known that equivariant cohomologies $H_G^*(X; \mathbb{Z}_2)$ and $H_G^*(X^G; \mathbb{Z}_2)$ are $H^*(BG; \mathbb{Z}_2)$ -modules; in particular, $H_G^*(X^G; \mathbb{Z}_2)$ is a free $H^*(BG; \mathbb{Z}_2)$ -module.

Suppose that M is a connected closed manifold and admits a G -action with M^G a non-empty finite fixed set. For the fibration $M \rightarrow M_G \rightarrow BG$, if the restriction to a typical fiber

$$H_G^*(M; \mathbb{Z}_2) \rightarrow H^*(M; \mathbb{Z}_2)$$

is an epimorphism, then M is called *totally non-homologous to zero* in M_G (see [B]). Under this condition, M is also called *G -equivariantly formal* (cf. [GKM]). Note that when the action group on M is replaced by a 2-torus $(\mathbb{Z}_2)^k$ with $k \geq 1$ and each

2000 Mathematics Subject Classification. 57S17, 55N91, 58J20

Key words and phrases. G -manifold, equivariant index, equivariant cohomology, analytic ring isomorphism.

Supported by grants from NSFC (No. 10371020 and No. 10671034).

component of M^K has dimension at most 1 for $K < (\mathbb{Z}_2)^k$ a corank-1 2-torus, if M is $(\mathbb{Z}_2)^k$ -equivariantly formal (i.e., the edge morphism $H_{(\mathbb{Z}_2)^k}^*(M; \mathbb{Z}_2) \longrightarrow H^*(M; \mathbb{Z}_2)$ is surjective), then there is a mod 2 GKM theory (corresponding to GKM theory, see [GKM], [GZ]), indicating that the $(\mathbb{Z}_2)^k$ -equivariant cohomology of M can be explicitly expressed in terms of its associated graph (Γ_M, α) (cf. [BGH], [L]). In particular, when $k = 1$, M is naturally restricted to have dimension at most 1, so this means that the GKM theory can be carried out only for at most 1-dimensional G -equivariantly formal manifolds. In this paper, we shall give explicit descriptions of equivariant cohomology rings of G -equivariantly formal manifolds at any dimension, and these descriptions are of algebra rather than combinatorics.

Let Λ_n denote the set of all n -dimensional connected closed G -manifolds with a non-empty finite fixed point set, each of which is G -equivariantly formal. Note that obviously Λ_1 contains a unique 1-dimensional closed manifold, i.e., a circle S^1 . Taking a M in Λ_n , by Conner and Floyd [CF], one knows that $|M^G|$ must be even. Let r be a positive integer, and write $\Lambda_n^{2r} = \{M \in \Lambda_n \mid |M^G| = 2r\}$. Then $\Lambda_n = \bigcup_{r \geq 1} \Lambda_n^{2r}$. Given a M in Λ_n , we know from [AP] and [B] that the following conditions are equivalent

- (1) M is G -equivariantly formal;
- (2) $|M^G| = \sum_{i=0}^n b_i$ where b_i is the i -th mod 2 Betti number of M ;
- (3) $H_G^*(M; \mathbb{Z}_2)$ is a free $H^*(BG; \mathbb{Z}_2)$ -module;
- (4) The inclusion $i : M^G \hookrightarrow M$ induces a monomorphism $i^* : H_G^*(M; \mathbb{Z}_2) \longrightarrow H_G^*(M^G; \mathbb{Z}_2)$.

If $|M^G| = 2r$, since $H_G^*(M^G; \mathbb{Z}_2) = \bigoplus_{p \in M^G} H_G^*(\{p\}; \mathbb{Z}_2)$ and the equivariant cohomology of a point is isomorphic to $H^*(BG; \mathbb{Z}_2) = \mathbb{Z}_2[t]$, we have that $H_G^*(M^G; \mathbb{Z}_2) \cong (\mathbb{Z}_2)^{2r}[t]$ is a polynomial ring (or algebra). Thus we obtain a monomorphism from $H_G^*(M; \mathbb{Z}_2)$ into $(\mathbb{Z}_2)^{2r}[t]$, also denoted by i^* , so $H_G^*(M; \mathbb{Z}_2)$ may be identified with a subring (or subalgebra) of $(\mathbb{Z}_2)^{2r}[t]$.

Using the equivariant index, we shall give an explicit description of $H_G^*(M; \mathbb{Z}_2)$ in $(\mathbb{Z}_2)^{2r}[t]$ (see Theorem 3.1). Then we consider the following questions:

- (Q1) If the equivariant cohomology rings of two G -manifolds in Λ_n are isomorphic, then can the isomorphism between them be explicitly expressed?
- (Q2) When are the equivariant cohomology rings of two G -manifolds in Λ_n isomorphic?
- (Q3) How many are there equivariant cohomology rings (up to isomorphism) of G -manifolds in Λ_n ?

We completely answer (Q1) and (Q2). An interesting thing is that we do not only find an explicit description for the isomorphism between equivariant cohomology rings of two G -manifolds in Λ_n , but such a description is also analytic (see Theorem 5.1), so that we may obtain a necessary and sufficient condition that the equivariant cohomology rings of such two G -manifolds are isomorphic in terms of algebra (see Theorem 5.2).

As for (Q3), the question is answered completely in the case $n = 2$. When $n = 3$, we find an upper bound of the number for the equivariant cohomology rings (up to isomorphism) of G -manifolds in Λ_3^{2r} (see Proposition 6.1).

The paper is organized as follows. In Section 2 we review the localization theorem and reformulate the equivariant index from the work of Allday and Puppe [AP]. In Section 3 we study the structure of equivariant cohomology of a G -manifold in Λ_n and obtain an explicit description in terms of algebra. Then we completely answer (Q2) and (Q3) in the case $n = 2$ in Section 4. In Section 5, we give an analytic description for the isomorphism between equivariant cohomology rings of two G -manifolds in Λ_n , so that we may obtain a necessary and sufficient condition that the equivariant cohomology rings of two G -manifolds in Λ_n are isomorphic. In section 6, we discuss the number of the equivariant cohomology rings (up to isomorphism) of G -manifolds in Λ_3^{2r} , and obtain an upper bound of the number.

The authors express their thanks to Shengzhi Xu for helpful conversation in the argument of Lemma 6.1. The authors also would like to express their gratitude to Professor Volker Puppe, who informed of us that there is an essential relationship between the equivariant cohomology rings and the coding theory (see [P]), and Theorem 3 in [KP] implies that the map g is surjective in the remark 8 of our paper.

2. LOCALIZATION THEOREM AND EQUIVARIANT INDEX

Suppose that M is an n -dimensional G -manifold with M^G a non-empty finite set. Let S be the subset of $H^*(BG; \mathbb{Z}_2)$ generated multiplicatively by nonzero elements in $H^1(BG; \mathbb{Z}_2)$. Then one has the following well-known localization theorem (see [AP], [H]).

Theorem 2.1 (Localization theorem).

$$S^{-1}i^* : S^{-1}H_G^*(M; \mathbb{Z}_2) \longrightarrow S^{-1}H_G^*(M^G; \mathbb{Z}_2)$$

is an isomorphism of $S^{-1}H^*(BG; \mathbb{Z}_2)$ -algebras, where i is the inclusion of from M^G into M .

Take an isolated point $p \in M^G$. Let i_p be the inclusion of from p into M , then one has the equivariant Gysin homomorphism

$$i_{p!} : H_G^*(\{p\}; \mathbb{Z}_2) \longrightarrow H_G^{*+n}(M; \mathbb{Z}_2).$$

On the other hand, one has also a natural induced homomorphism

$$i_p^* : H_G^*(M; \mathbb{Z}_2) \longrightarrow H_G^*(\{p\}; \mathbb{Z}_2)$$

and in particular, it is easy to check that $i^* = \bigoplus_{p \in M^G} i_p^*$. Furthermore, one knows that the equivariant Euler class at p is

$$\chi_G(p) = i_p^* i_{p!}(1_p) \in H_G^n(\{p\}; \mathbb{Z}_2) = H^n(BG; \mathbb{Z}_2) = \mathbb{Z}_2 t^n,$$

which is equal to that of the real G -representation at p , where $1_p \in H_G^*(\{p\}; \mathbb{Z}_2)$ is the identity and $\mathbb{Z}_2 t^n = \{at^n | a \in \mathbb{Z}_2\}$. Thus, we may write $\chi_G(p) = t^n$. Write $\theta_p = i_{p!}(1_p)$. Then $\theta_p \in H_G^n(M; \mathbb{Z}_2)$ and $i_p^*(\theta_p) = \chi_G(p)$.

Lemma 2.1. *All elements $\theta_p, p \in M^G$ are linearly independent over $H^*(BG; \mathbb{Z}_2)$.*

Proof. Let $\sum_{p \in M^G} l_p \theta_p = 0$, where $l_p \in H^*(BG; \mathbb{Z}_2)$. From [AP, Proposition 5.3.14(2)], one knows that $i_q^*(\theta_p) = 0$ for $q \neq p$ in M^G , so

$$i_q^*\left(\sum_{p \in M^G} l_p \theta_p\right) = \sum_{p \in M^G} l_p i_q^*(\theta_p) = l_q i_q^*(\theta_q) = l_q \chi_G(q) = 0.$$

Since $\chi_G(q) = t^n$ is a unit in $S^{-1}H_G^*(\{q\}; \mathbb{Z}_2) \cong S^{-1}H^*(BG; \mathbb{Z}_2)$, one has $l_q = 0$. $\square$

Lemma 2.2. *Let $\alpha \in S^{-1}H_G^*(M; \mathbb{Z}_2)$. Then*

$$\alpha = \sum_{p \in M^G} \frac{f_p \theta_p}{t^n}$$

where $f_p = S^{-1}i_p^*(\alpha) \in S^{-1}H^*(BG; \mathbb{Z}_2)$.

Proof. By [AP, Proposition 5.3.18(1)], one has that

$$\alpha = \sum_{p \in M^G} S^{-1}i_{p!}(S^{-1}i_p^*(\alpha)/\chi_G(p)).$$

Since $f_p = S^{-1}i_p^*(\alpha) \in S^{-1}H^*(BG; \mathbb{Z}_2)$, one has that $\frac{f_p}{\chi_G(p)} = \frac{f_p}{t^n} \in S^{-1}H^*(BG; \mathbb{Z}_2)$. Since $S^{-1}i_{p!}$ is a $S^{-1}H^*(BG; \mathbb{Z}_2)$ -algebra homomorphism, one has

$$S^{-1}i_{p!}(S^{-1}i_p^*(\alpha)/\chi_G(p)) = S^{-1}i_{p!}\left(\frac{f_p}{t^n}\right) = \frac{f_p}{t^n} S^{-1}i_{p!}(1_p) = \frac{f_p}{t^n} i_{p!}(1_p) = \frac{f_p \theta_p}{t^n}$$

so $\alpha = \sum_{p \in M^G} \frac{f_p \theta_p}{t^n}$. $\square$

Remark 1. (i) By Lemma 2.1, one sees from the formula of Lemma 2.2 that $\{\frac{\theta_p}{t^n} | p \in M^G\}$ forms a basis of $S^{-1}H_G^*(M; \mathbb{Z}_2)$ as a $S^{-1}H^*(BG; \mathbb{Z}_2)$ -algebra.

(ii) In some sense, the formula $\alpha = \sum_{p \in M^G} \frac{f_p \theta_p}{t^n}$ explicitly indicates the isomorphism $S^{-1}i^* : S^{-1}H_G^*(M; \mathbb{Z}_2) \longrightarrow S^{-1}H_G^*(M^G; \mathbb{Z}_2)$ in Theorem 2.1, which is given by mapping $\alpha = \sum_{p \in M^G} \frac{f_p \theta_p}{t^n}$ to $\bigoplus_{p \in M^G} \frac{f_p}{t^n}$.

The equivariant Gysin homomorphism of collapsing M to a point gives the G -index of M , i.e.,

$$\text{Ind}_G : H_G^*(M; \mathbb{Z}_2) \longrightarrow H^{*-n}(BG; \mathbb{Z}_2).$$

Theorem 2.2. *For any $\alpha \in S^{-1}H_G^*(M; \mathbb{Z}_2)$,*

$$S^{-1}\text{Ind}_G(\alpha) = \sum_{p \in M^G} \frac{f_p}{t^n}$$

where $f_p = S^{-1}i_p^*(\alpha) \in S^{-1}H^*(BG; \mathbb{Z}_2)$. In particular, if $\alpha \in H_G^*(M; \mathbb{Z}_2)$, then $f_p = i_p^*(\alpha) \in H^*(BG; \mathbb{Z}_2)$ and

$$(1) \quad \text{Ind}_G(\alpha) = \sum_{p \in M^G} \frac{f_p}{t^n} \in H^*(BG; \mathbb{Z}_2) = \mathbb{Z}_2[t].$$

Proof. By [AP, Lemma 5.3.19], one has that $\text{Ind}_G(\theta_p) = 1_p$, so by Lemma 2.2

$$S^{-1}\text{Ind}_G(\alpha) = \sum_{p \in M^G} \frac{f_p S^{-1}\text{Ind}_G(\theta_p)}{t^n} = \sum_{p \in M^G} \frac{f_p \cdot 1_p}{t^n} = \sum_{p \in M^G} \frac{f_p}{t^n}.$$

The last part of Theorem 2.2 follows immediately since $H^*(BG; \mathbb{Z}_2) \longrightarrow S^{-1}H^*(BG; \mathbb{Z}_2)$ is injective. $\square$

Remark 2. a) It should be pointed out that all arguments in this section can still be carried out if the action group G is a 2-torus $(\mathbb{Z}_2)^k$ of rank $k > 1$. In this case, $H^*(BG; \mathbb{Z}_2)$ is a polynomial algebra $\mathbb{Z}_2[t_1, \dots, t_k]$ where the t_i 's are one-dimensional generators in $H^1(BG; \mathbb{Z}_2)$, so that the formula (1) becomes

$$(2) \quad \text{Ind}_G(\alpha) = \sum_{p \in M^G} \frac{f_p}{\chi_G(p)} \in \mathbb{Z}_2[t_1, \dots, t_k]$$

where $\chi_G(p)$ is a polynomial of degree n in $\mathbb{Z}_2[t_1, \dots, t_k]$. Note that related results can also be found in [D] and [KS].

b) The formula (2) is an analogue of the Atiyah-Bott-Berlin-Vergne formula for the case $G = T$ (i.e., a torus), see [AB2] and [BV].

3. EQUIVARIANT COHOMOLOGY STRUCTURE

In this section, our task is to study the structures of equivariant cohomology rings of G -manifolds in Λ_n .

Lemma 3.1. *Let $M \in \Lambda_n^{2r}$. Then*

$$\dim_{\mathbb{Z}_2} H_G^i(M; \mathbb{Z}_2) = \begin{cases} \sum_{j=0}^i b_j & \text{if } i \leq n-1 \\ 2r & \text{if } i \geq n. \end{cases}$$

Proof. Let

$$P_s(M_G) = \sum_{i=0}^{\infty} \dim_{\mathbb{Z}_2} H_G^i(M; \mathbb{Z}_2) s^i$$

be the equivariant Poincaré polynomial of $H_G^*(X; \mathbb{Z}_2)$. Since $H_G^*(M; \mathbb{Z}_2)$ is a free $H^*(BG; \mathbb{Z}_2)$ -module, one has that $H_G^*(M; \mathbb{Z}_2) = H^*(M; \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} H^*(BG; \mathbb{Z}_2)$ so

$$P_s(M_G) = \sum_{i=0}^{\infty} \dim_{\mathbb{Z}_2} H_G^i(M; \mathbb{Z}_2) s^i = \frac{1}{1-s} \sum_{i=0}^n \dim_{\mathbb{Z}_2} H^i(M; \mathbb{Z}_2) s^i.$$

Write $b_i = \dim_{\mathbb{Z}_2} H^i(M; \mathbb{Z}_2)$. Note that $b_i = b_{n-i}$ by Poincaré duality and $b_0 = b_n = 1$ since M is connected. Then

$$\begin{aligned} P_s(M_G) &= \sum_{i=0}^{\infty} \dim_{\mathbb{Z}_2} H_G^i(M; \mathbb{Z}_2) s^i = \frac{1}{1-s} \sum_{i=0}^n \dim_{\mathbb{Z}_2} H^i(M; \mathbb{Z}_2) s^i \\ &= b_0 + (b_0 + b_1)s + \cdots + (b_0 + b_1 + \cdots + b_{n-1})s^{n-1} + (b_0 + b_1 + \cdots + b_n)(s^n + \cdots) \\ &= b_0 + (b_0 + b_1)s + \cdots + (b_0 + b_1 + \cdots + b_{n-1})s^{n-1} + 2r(s^n + \cdots). \end{aligned}$$

The lemma then follows from this. $\square$

Let $x = (x_1, \dots, x_{2r})$ and $y = (y_1, \dots, y_{2r})$ be two vectors in $(\mathbb{Z}_2)^{2r}$. Define $x \circ y$ by

$$x \circ y = (x_1 y_1, \dots, x_{2r} y_{2r}).$$

Then $(\mathbb{Z}_2)^{2r}$ forms a commutative ring with respect to two operations $+$ and $\circ$. Let

$$\mathcal{V}_{2r} = \left\{ x = (x_1, \dots, x_{2r})^\top \in (\mathbb{Z}_2)^{2r} \mid |x| = \sum_{i=1}^{2r} x_i = 0 \right\}.$$

Then it is easy to see that $\mathcal{V}_{2r}$ is a $(2r-1)$ -dimensional subspace of $(\mathbb{Z}_2)^{2r}$, and there is only such a subspace in $(\mathbb{Z}_2)^{2r}$. However, generally the operation $\circ$ in $\mathcal{V}_{2r}$ is obviously not closed.

Given a $M \in \Lambda_n^{2r}$, one then has that the inclusion $i : M^G \hookrightarrow M$ induces a monomorphism

$$i^* : H_G^*(M; \mathbb{Z}_2) \longrightarrow (\mathbb{Z}_2)^{2r}[t].$$

By Lemma 3.1, there are subspaces V_i^M with $\dim V_i^M = \sum_{j=0}^i b_j$ ($i = 0, \dots, n-1$) of $(\mathbb{Z}_2)^{2r}$ such that

$$H_G^i(M; \mathbb{Z}_2) \cong i^*(H_G^i(M; \mathbb{Z}_2)) = \begin{cases} V_i^M t^i & \text{if } i \leq n-1 \\ (\mathbb{Z}_2)^{2r} t^i & \text{if } i \geq n \end{cases}$$

where $V_i^M t^i = \{vt^i \mid v \in V_i^M\}$.

Lemma 3.2. *There are the following properties:*

- (1) $\mathbb{Z}_2 \cong V_0^M \subset V_1^M \subset \cdots \subset V_{n-2}^M \subset V_{n-1}^M = \mathcal{V}_{2r}$, where V_0^M is generated by $(1, \dots, 1)^\top \in (\mathbb{Z}_2)^{2r}$;
- (2) For $d = \sum_{i=0}^{n-1} id_i < n$ with each $d_i \geq 0$, $v_{\omega_{d_0}} \circ \cdots \circ v_{\omega_{d_{n-1}}} \in V_d^M$, where $v_{\omega_{d_i}} = v_1^{(i)} \circ \cdots \circ v_{d_i}^{(i)}$ with each $v_j^{(i)} \in V_i^M$.

Proof. For an element $\alpha \in H_G^*(M; \mathbb{Z}_2)$ of degree d , one has that $i^*(\alpha) = vt^d$ where $v \in (\mathbb{Z}_2)^{2r}$. Since $i^* = \bigoplus_{p \in M^G} i_p^*$, by Theorem 2.2 one has that

$$\text{Ind}_G(\alpha) = \sum_{p \in M^G} \frac{i_p^*(\alpha)}{t^n} = \frac{1}{t^n} \sum_{p \in M^G} i_p^*(\alpha) = |v|t^{d-n} \in \mathbb{Z}_2[t]$$

so if $d < n$, then $|v|$ must be zero. This means that for each $i < n$, V_i^M is a subspace of $\mathcal{V}_{2r}$ and $V_{n-1}^M = \mathcal{V}_{2r}$ is obvious since $\dim V_{n-1}^M = 2r - 1$. In particular, when $\alpha = 1$ is the identity of $H_G^*(M; \mathbb{Z}_2)$, $i^*(1) = \bigoplus_{p \in M^G} i_p^*(1) = (1, \dots, 1)^\top \in (\mathbb{Z}_2)^{2r}$. Thus, $V_0^M \cong \mathbb{Z}_2$ is generated by $(1, \dots, 1)^\top \in (\mathbb{Z}_2)^{2r}$ since $\dim V_0^M = b_0 = 1$. Since $H_G^*(M; \mathbb{Z}_2) = H^*(M; \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} H^*(BG; \mathbb{Z}_2)$, one has that $(1, \dots, 1)^\top t \in i^*(H_G^1(M; \mathbb{Z}_2))$. Thus, for any $v \in V_i^M$ with $i < n - 1$, $(v t^i) \circ [(1, \dots, 1)^\top t] = v t^{i+1} \in V_{i+1}^M t^{i+1}$, so one has that $v \in V_{i+1}^M$. This completes the proof of Lemma 3.2(1).

As for the proof of Lemma 3.2(2), for each $v_j^{(i)} \in V_i^M$, since $i^* : H_G^*(M; \mathbb{Z}_2) \longrightarrow (\mathbb{Z}_2)^{2r}[t]$ is injective, there is a class $\alpha_j^{(i)}$ of degree i in $H_G^*(M; \mathbb{Z}_2)$ such that $i^*(\alpha_j^{(i)}) = v_j^{(i)} t^i$. Since $i^* = \bigoplus_{p \in M^G} i_p^*$ is also a ring homomorphism, one has that

$$i^*\left(\prod_{i=0}^{n-1} \prod_{j=1}^{d_i} \alpha_j^{(i)}\right) = \bigoplus_{p \in M^G} i_p^*\left(\prod_{i=0}^{n-1} \prod_{j=1}^{d_i} \alpha_j^{(i)}\right) = \bigoplus_{p \in M^G} \prod_{i=0}^{n-1} \prod_{j=1}^{d_i} i_p^*(\alpha_j^{(i)}) = v_{\omega_{d_0}} \circ \dots \circ v_{\omega_{d_{n-1}}} t^d$$

so $v_{\omega_{d_0}} \circ \dots \circ v_{\omega_{d_{n-1}}} \in V_d^M$. $\square$

Remark 3. An easy observation shows that the properties (1) and (2) of Lemma 3.2 exactly give a subring structure of

$$\mathcal{R}_M = V_0^M + V_1^M t + \dots + V_{n-1}^M t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \dots)$$

in $(\mathbb{Z}_2)^{2r}[t]$.

Combining Lemmas 3.1, 3.2, and Remark 3 one has

Theorem 3.1. *Let $M \in \Lambda_n^{2r}$. Then there are subspaces V_i^M with $\dim V_i^M = \sum_{j=0}^i b_j$ ($i = 0, \dots, n - 1$) of $\mathcal{V}_{2r}$ such that $H_G^*(M; \mathbb{Z}_2)$ is isomorphic to the graded ring*

$$\mathcal{R}_M = V_0^M + V_1^M t + \dots + V_{n-2}^M t^{n-2} + V_{n-1}^M t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + t^{n+1} + \dots)$$

where the ring structure of $\mathcal{R}_M$ is given by

- (a) $\mathbb{Z}_2 \cong V_0^M \subset V_1^M \subset \dots \subset V_{n-2}^M \subset V_{n-1}^M = \mathcal{V}_{2r}$, where V_0^M is generated by $(1, \dots, 1)^\top \in (\mathbb{Z}_2)^{2r}$;
- (b) For $d = \sum_{i=0}^{n-1} i d_i < n$ with each $d_i \geq 0$, $v_{\omega_{d_0}} \circ \dots \circ v_{\omega_{d_{n-1}}} \in V_d^M$, where $v_{\omega_{d_i}} = v_1^{(i)} \circ \dots \circ v_{d_i}^{(i)}$ with each $v_j^{(i)} \in V_i^M$.

Remark 4. Since $H_G^*(M; \mathbb{Z}_2)$ is also a free $H^*(BG; \mathbb{Z}_2)$ -module, one has that

$$\mathcal{R}_M = V_0^M + V_1^M t + \dots + V_{n-2}^M t^{n-2} + V_{n-1}^M t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + t^{n+1} + \dots)$$

is a free $\mathbb{Z}_2[t]$ -module, too.

Next, let us determine the largest-dimensional space V in $\{V_i^M | 0 \leq i \leq n - 1\}$ with the property that $u \circ v \in \mathcal{V}_{2r}$ for $u, v \in V$.

Given a vector $v \in \mathcal{V}_{2r}$, set

$$\mathcal{V}(v) := \{x \in \mathcal{V}_{2r} | x \circ v \in \mathcal{V}_{2r}\}.$$

Then it is easy to see that $\mathcal{V}(v)$ is a linear subspace of $\mathcal{V}_{2r}$, and $\mathcal{V}(\underline{1} + v) = \mathcal{V}(v)$, and $\mathcal{V}(v) = \mathcal{V}_{2r} \iff v = \underline{0}$ or $\underline{1}$ where $\underline{0} = (0, \dots, 0)^\top$ and $\underline{1} = (1, \dots, 1)^\top$.

Lemma 3.3. *Let $v \in \mathcal{V}_{2r}$ with $v \neq \underline{0}, \underline{1}$. Then $\dim \mathcal{V}(v) = 2r - 2$.*

Proof. Let $\mathcal{V}_1(v) = \{x \in \mathcal{V}_{2r} | v \circ x = x\}$ and $\mathcal{V}_2(v) = \{x \in \mathcal{V}_{2r} | v \circ x = \underline{0}\}$. Obviously, they are subspace of $\mathcal{V}_{2r}$ and $\mathcal{V}_1(v) \cap \mathcal{V}_2(v) = \{\underline{0}\}$. For any $x \in \mathcal{V}(v)$, $x = v \circ x + (v \circ x + x)$. Since $v \circ x \in \mathcal{V}_1(v)$ and $v \circ x + x \in \mathcal{V}_2(v)$, one has that $\mathcal{V}(v) \subset \mathcal{V}_1(v) \oplus \mathcal{V}_2(v)$. However, obviously $\mathcal{V}_1(v) \oplus \mathcal{V}_2(v) \subset \mathcal{V}(v)$. Thus, $\mathcal{V}(v) = \mathcal{V}_1(v) \oplus \mathcal{V}_2(v)$.

By $\#v$ one denotes the number of nonzero elements in v . Let $\#v = m$. Then $0 < m < 2r$ since $v \neq \underline{0}, \underline{1}$, and m is even since $v \in \mathcal{V}_{2r}$. An easy observation shows that $\dim \mathcal{V}_1(v) = m - 1$ and $\dim \mathcal{V}_2(v) = (2r - m) - 1$. Thus,

$$\dim \mathcal{V}(v) = \dim \mathcal{V}_1(v) + \dim \mathcal{V}_2(v) = m - 1 + (2r - m) - 1 = 2r - 2.$$

□

Lemma 3.4. *Let $u, v \in \mathcal{V}_{2r}$. Then*

$$\mathcal{V}(u + v) = [\mathcal{V}(u) \cap \mathcal{V}(v)] \cup [\mathcal{V}_{2r} \setminus (\mathcal{V}(u) \cup \mathcal{V}(v))].$$

Proof. For any $x \in \mathcal{V}(u + v)$, one has that $x \circ (u + v) = x \circ u + x \circ v \in \mathcal{V}_{2r}$ so $\#(x \circ u + x \circ v)$ is even. Since $\#(a + b) = \#a + \#b - 2\#(a \circ b)$ for any $a, b \in (\mathbb{Z}_2)^{2r}$, one has that both $\#(x \circ u)$ and $\#(x \circ v)$ are even or odd. Thus

$$\begin{aligned} & \mathcal{V}(u + v) \\ &= \{x \in \mathcal{V}_{2r} | x \circ u + x \circ v \in \mathcal{V}_{2r}\} \\ &= \{x \in \mathcal{V}_{2r} | \#(x \circ u) \text{ and } \#(x \circ v) \text{ are even}\} \cup \{x \in \mathcal{V}_{2r} | \#(x \circ u) \text{ and } \#(x \circ v) \text{ are odd}\} \\ &= [\mathcal{V}(u) \cap \mathcal{V}(v)] \cup [\mathcal{V}_{2r} \setminus (\mathcal{V}(u) \cup \mathcal{V}(v))]. \end{aligned}$$

□

Let $v_1, \dots, v_k \in \mathcal{V}_{2r}$, and let $\mathcal{V}(v_1, \dots, v_k)$ denote $\mathcal{V}(v_1) \cap \dots \cap \mathcal{V}(v_k)$.

Proposition 3.1. *Suppose that $\underline{1}, v_1, \dots, v_k \in \mathcal{V}_{2r}$ are linearly independent. Then*

$$\dim \mathcal{V}(v_1, \dots, v_k) = 2r - 1 - k.$$

Proof. One uses induction on k . From Lemma 3.3 one knows that the case $k = 1$ holds. If $k \leq m$, suppose inductively that Proposition 3.1 holds. Consider the case $k = m + 1$. Since $\mathcal{V}(v_1, \dots, v_{m+1}) \subset \mathcal{V}(v_1, \dots, v_m)$, one has $\dim \mathcal{V}(v_1, \dots, v_{m+1}) \leq \dim \mathcal{V}(v_1, \dots, v_m)$.

If $\dim \mathcal{V}(v_1, \dots, v_{m+1}) = 2r - 1 - m$, then $\mathcal{V}(v_1, \dots, v_{m+1}) = \mathcal{V}(v_1, \dots, v_m)$. One claims that this is impossible. If so, by induction, $\dim \mathcal{V}(v_2, \dots, v_{m+1}) = 2r - 1 - m$ so

$$(3) \quad \mathcal{V}(v_2, \dots, v_{m+1}) = \mathcal{V}(v_1, \dots, v_m)$$

since $\mathcal{V}(v_1, \dots, v_m) = \mathcal{V}(v_1, \dots, v_{m+1}) \subset \mathcal{V}(v_2, \dots, v_{m+1})$. In a similar way, one has also that

$$(4) \quad \mathcal{V}(v_1, v_3, \dots, v_{m+1}) = \mathcal{V}(v_1, \dots, v_m).$$

By Lemma 3.4,

$$\begin{aligned} \mathcal{V}(v_1 + v_2) \cap \mathcal{V}(v_1) &= \{[\mathcal{V}(v_1) \cap \mathcal{V}(v_2)] \cup [\mathcal{V}_{2r} \setminus (\mathcal{V}(v_1) \cup \mathcal{V}(v_2))]\} \cap \mathcal{V}(v_1) \\ &= [\mathcal{V}(v_1) \cap \mathcal{V}(v_2)] \cap \emptyset \\ &= \mathcal{V}(v_1) \cap \mathcal{V}(v_2). \end{aligned}$$

Similarly, one also has that $\mathcal{V}(v_1 + v_2) \cap \mathcal{V}(v_2) = \mathcal{V}(v_1) \cap \mathcal{V}(v_2)$. Thus

$$\begin{aligned} &\mathcal{V}(v_1 + v_2, v_3, \dots, v_{m+1}) \cap \mathcal{V}(v_1, \dots, v_m) \\ &= \{\mathcal{V}(v_1 + v_2) \cap \mathcal{V}(v_3, \dots, v_{m+1})\} \cap \{\mathcal{V}(v_1) \cap \mathcal{V}(v_2) \cap \mathcal{V}(v_3, \dots, v_m)\} \\ &= \mathcal{V}(v_1, \dots, v_m) \end{aligned}$$

so

$$\mathcal{V}(v_1 + v_2, v_3, \dots, v_{m+1}) = \mathcal{V}(v_1, \dots, v_m).$$

On the other hand,

$$\begin{aligned} &\mathcal{V}(v_1 + v_2, v_3, \dots, v_{m+1}) (= \mathcal{V}(v_1, \dots, v_m) = \mathcal{V}(v_1, \dots, v_{m+1})) \\ &= \mathcal{V}(v_1 + v_2) \cap \mathcal{V}(v_3, \dots, v_{m+1}) \\ &= \{[\mathcal{V}(v_1) \cap \mathcal{V}(v_2)] \cup [\mathcal{V}_{2r} \setminus (\mathcal{V}(v_1) \cup \mathcal{V}(v_2))]\} \cap \mathcal{V}(v_3, \dots, v_{m+1}) \text{ by Lemma 3.4} \\ &= \mathcal{V}(v_1, \dots, v_{m+1}) \cup \{[\mathcal{V}_{2r} \setminus (\mathcal{V}(v_1) \cup \mathcal{V}(v_2))] \cap \mathcal{V}(v_3, \dots, v_{m+1})\}. \end{aligned}$$

This implies that

$$(5) \quad [\mathcal{V}_{2r} \setminus (\mathcal{V}(v_1) \cup \mathcal{V}(v_2))] \cap \mathcal{V}(v_3, \dots, v_{m+1}) = \emptyset.$$

Combining the formulae (3), (4) and (5), one obtains that

$$\begin{aligned} \mathcal{V}(v_1, \dots, v_m) &= [\mathcal{V}(v_1) \cup \mathcal{V}(v_2)] \cap \mathcal{V}(v_3, \dots, v_{m+1}) \text{ by (3) and (4)} \\ &= \{\mathcal{V}(v_1) \cup \mathcal{V}(v_2) \cup [\mathcal{V}_{2r} \setminus (\mathcal{V}(v_1) \cup \mathcal{V}(v_2))]\} \cap \mathcal{V}(v_3, \dots, v_{m+1}) \text{ by (5)} \\ &= \mathcal{V}_{2r} \cap \mathcal{V}(v_3, \dots, v_{m+1}) \\ &= \mathcal{V}(v_3, \dots, v_{m+1}) \end{aligned}$$

so $\mathcal{V}(v_1, \dots, v_m) = \mathcal{V}(v_3, \dots, v_{m+1})$. However, by induction, $\dim \mathcal{V}(v_1, \dots, v_m) = 2r - 1 - m$ but $\dim \mathcal{V}(v_3, \dots, v_{m+1}) = 2r - 1 - (m - 1)$. This is a contradiction.

Therefore, $\dim \mathcal{V}(v_1, \dots, v_{m+1}) < \dim \mathcal{V}(v_1, \dots, v_m)$.

Now let $\{x_1, \dots, x_{2r-1-m}\}$ be a basis of $\mathcal{V}(v_1, \dots, v_m)$. Since

$$\mathcal{V}(v_1, \dots, v_{m+1}) \subsetneq \mathcal{V}(v_1, \dots, v_m)$$

there must exist at least one element x in $\{x_1, \dots, x_{2r-1-m}\}$ such that $x \notin \mathcal{V}(v_1, \dots, v_{m+1})$. With no loss, one may assume that $x_1 \notin \mathcal{V}(v_1, \dots, v_{m+1})$. Then $x_1 \circ v_{m+1} \notin \mathcal{V}_{2r}$. If there is also another $x_i (i \neq 1)$ in $\{x_1, x_2, \dots, x_{2r-1-m}\}$ such that $x_i \notin \mathcal{V}(v_1, \dots, v_{m+1})$,

then $x_i \circ v_{m+1} \notin \mathcal{V}_{2r}$. However, one knows from the proof of Lemma 3.4 that $x_1 \circ v_{m+1} + x_i \circ v_{m+1} \in \mathcal{V}_{2r}$ so $x_1 + x_i \in \mathcal{V}(v_{m+1})$ and $x_1 + x_i \in \mathcal{V}(v_1, \dots, v_{m+1})$. In this case, $\{x_1, \dots, x_{i-1}, x_1 + x_i, x_{i+1}, \dots, x_{2r-1-m}\}$ is still a basis of $\mathcal{V}(v_1, \dots, v_m)$ but $x_1 + x_i \in \mathcal{V}(v_1, \dots, v_{m+1})$. One can use this way to further modify the basis $\{x_1, \dots, x_{i-1}, x_1 + x_i, x_{i+1}, \dots, x_{2r-1-m}\}$ into a basis $\{x_1, x'_2, \dots, x'_{2r-1-m}\}$, such that $x'_2, \dots, x'_{2r-1-m} \in \mathcal{V}(v_1, \dots, v_{m+1})$ except for $x_1 \notin \mathcal{V}(v_1, \dots, v_{m+1})$. Thus, $\dim \mathcal{V}(v_1, \dots, v_{m+1}) = 2r - 1 - (m + 1)$. This completes the induction and the proof of Proposition 3.1. $\square$

Corollary 3.1. *Suppose that $\underline{1}, v_1, \dots, v_k \in \mathcal{V}_{2r}$ are linearly independent with $v_i \circ v_j \in \mathcal{V}_{2r}$ for any $i, j \in \{1, \dots, k\}$. Then $k \leq r - 1$.*

Proof. Since $v_i \circ v_j \in \mathcal{V}_{2r}$ for any $i, j \in \{1, \dots, k\}$,

$$\text{Span}\{\underline{1}, v_1, \dots, v_k\} \subset \mathcal{V}(v_1, \dots, v_k).$$

So $k + 1 \leq 2r - 1 - k$, i.e., $k \leq r - 1$. $\square$

Corollary 3.2. *The largest-dimensional space of preserving the operation $\circ$ closed in $\mathcal{V}_{2r}$ of $\{V_i^M | 0 \leq i \leq n - 1\}$ is $V_{[\frac{n}{2}]}^M$ with dimension r if n is odd, and $V_{\frac{n}{2}-1}^M$ with dimension $< r$ if n is even but $b_{\frac{n}{2}} \neq 0$, and $V_{\frac{n}{2}-1}^M = V_{\frac{n}{2}}^M$ with dimension r if n is even and $b_{\frac{n}{2}} = 0$.*

Proof. Since $b_0 + b_1 + \dots + b_{n-1} + b_n = 2r$ and $b_i = b_{n-i}$ (note $b_0 = b_n = 1$), if n is odd, then $b_0 + b_1 + \dots + b_{[\frac{n}{2}]} = r$ so $\dim V_{[\frac{n}{2}]}^M = r$. Since $\underline{1} \in V_i^M$ for each i , by Corollary 3.1, the largest-dimensional space of preserving the operation $\circ$ closed in $\mathcal{V}_{2r}$ of $\{V_i^M | 0 \leq i \leq n - 1\}$ must be $V_{[\frac{n}{2}]}^M$. If n is even and $b_{\frac{n}{2}} \neq 0$, then $b_0 + b_1 + \dots + b_{\frac{n}{2}-1} < r$ but $b_0 + b_1 + \dots + b_{\frac{n}{2}} > r$ so the required largest-dimensional subspace is $V_{\frac{n}{2}-1}^M$ with dimension $r - \frac{b_{\frac{n}{2}}}{2} < r$. If n is even and $b_{\frac{n}{2}} = 0$, then

$$b_0 + b_1 + \dots + b_{\frac{n}{2}-1} = b_0 + b_1 + \dots + b_{\frac{n}{2}} = r$$

so the desired result holds. $\square$

4. 2-DIMENSIONAL CASE

Let $M \in \Lambda_2^{2r}$. By Theorem 3.1, we know that

$$H_G^*(M; \mathbb{Z}_2) \cong V_0^M + \mathcal{V}_{2r}t + (\mathbb{Z}_2)^{2r}(t^2 + t^3 + \dots).$$

This means that the ring structure of $H_G^*(M; \mathbb{Z}_2)$ only depends upon the number $2r = |M^G|$. Thus we have

Proposition 4.1. *Let $M_1, M_2 \in \Lambda_2$. Then $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ are isomorphic if and only if $|M_1^G| = |M_2^G|$.*

An easy observation shows that for an orientable connected closed surface Σ_g with genus $g \geq 0$, Σ_g must admit a G -action such that $|\Sigma_g^G| = 2(g + 1)$. Thus, for each $r \geq 1$, Λ_2^{2r} is non-empty.

Remark 5. However, one knows from [B] that $\mathbb{R}P^2$ never admits a G -action such that the fixed point set is a finite set so $\mathbb{R}P^2$ doesn't belong to Λ_2 . Actually, generally each non-orientable connected closed surface S_g with genus g odd must not belong to Λ_2 . This is because the sum of all mod 2 Betti numbers of S_g is $2 + g$, which is odd.

As a consequence of Proposition 4.1, one has

Corollary 4.1. *For each positive integer r , all G -manifolds in Λ_2^{2r} determine a unique equivariant cohomology up to isomorphism.*

Remark 6. For each $M \in \Lambda_2^{2r}$, its equivariant cohomology $H_G^*(M; \mathbb{Z}_2)$ may be expressed in a simpler way. Since $(1, \dots, 1)^\top \in V_0^M \cong \mathbb{Z}_2$ and $|v| = 0$ for $v \in \mathcal{V}_{2r}$, one has that

$$H_G^*(M; \mathbb{Z}_2) \cong \left\{ \alpha = (\alpha_1, \dots, \alpha_{2r}) \in (\mathbb{Z}_2)^{2r}[t] \mid \begin{cases} \alpha_1 = \dots = \alpha_{2r} & \text{if } \deg \alpha = 0 \\ \sum_{i=1}^{2r} \alpha_i = 0 & \text{if } \deg \alpha = 1 \end{cases} \right\}.$$

Compare with [GH, Proposition 3.1], Goldin and Holm gave a description for the equivariant cohomology of a compact connected symplectic 4-dimensional manifold with an effective Hamiltonian S^1 -action with a finite fixed set, so that they computed the equivariant cohomology of certain manifolds with a Hamiltonian action of a torus T (see [GH, Theorem 2]). Similarly to the argument in [GH], using this description of $H_G^*(M; \mathbb{Z}_2)$ and the main result of Chang and Skjelbred [CS], one may give an explicit description in $\mathbb{Z}_2[t_1, \dots, t_k]$ of the equivariant cohomology of a closed $(\mathbb{Z}_2)^k$ -manifold N with the following conditions:

- (1) The fixed point set is finite;
- (2) The equivariant cohomology of N is a free $H^*(B(\mathbb{Z}_2)^k; \mathbb{Z}_2)$ -module;
- (3) For $K < (\mathbb{Z}_2)^k$ a corank-1 2-torus, each component of N^K has dimension at most 2.

We would like to leave it to readers as an exercise. For this description in $\mathbb{Z}_2[t_1, \dots, t_k]$ of $H_{(\mathbb{Z}_2)^k}^*(N; \mathbb{Z}_2)$, actually the above restriction condition (3) for each component of N^K is the best possible since generally there can be different equivariant cohomology structures for G -manifolds in Λ_n^{2r} when $n \geq 3$ (see, e.g., Section 6 of this paper).

5. AN ANALYTIC DESCRIPTION OF RING ISOMORPHISMS AND A NECESSARY AND SUFFICIENT CONDITION

The purpose of this section is to give an analytic description for the isomorphism between equivariant cohomology rings of two G -manifolds in Λ_n and to show a necessary and sufficient condition that the equivariant cohomology rings of two G -manifolds in Λ_n are isomorphic.

Let

$$\mathcal{W}_{2r} = \{ \sigma \in \text{Aut} \mathcal{V}_{2r} \subset \text{GL}(2r, \mathbb{Z}_2) \mid \sigma(x \circ y) = \sigma(x) \circ \sigma(y) \text{ for any } x, y \in \mathcal{V}_{2r} \}.$$

Given two σ, τ in $\mathcal{W}_{2r}$, it is easy to check that $\sigma\tau \in \mathcal{W}_{2r}$. Thus one has that $\mathcal{W}_{2r}$ is a subgroup of $\text{Aut} \mathcal{V}_{2r}$.

Lemma 5.1. $\mathcal{W}_{2r}$ is the Weyl subgroup of $\mathrm{GL}(2r, \mathbb{Z}_2)$.

Proof. By [AB1], it suffices to prove that $\mathcal{W}_{2r}$ is isomorphic to the symmetric group $\mathcal{S}_{2r}$ of rank $2r$. Let $\sigma = (a_{ij})_{2r \times 2r}$ be an element of $\mathcal{W}_{2r} \subset \mathrm{GL}(2r, \mathbb{Z}_2)$. Since σ is an automorphism of $\mathcal{V}_{2r}$, there exists a vector $\underline{x} \in \mathcal{V}_{2r}$ such that $\sigma(\underline{x}) = \underline{1}$ where $\underline{1} = (1, \underbrace{\dots}_{2r}, 1)^\top$. Since $\underline{1} \circ x = x$ for any $x \in \mathcal{V}_{2r}$, one has that $\underline{1} = \sigma(\underline{x}) = \sigma(\underline{1} \circ \underline{x}) = \sigma(\underline{1}) \circ \sigma(\underline{x}) = \sigma(\underline{1})$ so

$$(6) \quad \sum_{j=1}^{2r} a_{ij} = 1, \text{ for } i = 1, 2, \dots, 2r.$$

On the other hand, for any $x = (x_1, \dots, x_{2r})^\top$ and $y = (y_1, \dots, y_{2r})^\top$ in $\mathcal{V}_{2r}$, one has that $\sigma(x \circ y) = \sigma(x) \circ \sigma(y)$ so

$$(7) \quad \sum_{l=1}^{2r} a_{il} x_l y_l = \left(\sum_{j=1}^{2r} a_{ij} x_j \right) \left(\sum_{k=1}^{2r} a_{ik} y_k \right), \text{ for } i = 1, 2, \dots, 2r.$$

From (6) one knows that for each i , the number $q(i)$ of nonzero elements in $a_{i1}, \dots, a_{i2r}$ is odd.

Now let us show that for each i , $q(i)$ actually must be 1. Taking an i , without loss of generality one may assume that $a_{i1} = \dots = a_{iq(i)} = 1$, and $a_{i(q(i)+1)} = \dots = a_{i2r} = 0$. Then from (7) one has

$$(8) \quad \left(\sum_{j=1}^{q(i)} x_j \right) \left(\sum_{k=1}^{q(i)} y_k \right) + \sum_{l=1}^{q(i)} x_l y_l = 0.$$

If $q(i) > 1$, taking x with $x_1 = x_{q(i)} = 1$ and $x_j = 0$ for $j \neq 1, q(i)$ and y with $y_2 = y_{q(i)} = 1$ and $y_k = 0$ for $k \neq 2, q(i)$, the left side of (8) then becomes 1, but this is impossible. Thus $q(i) = 1$.

Since $q(i) = 1$ for each i , this means that σ is actually obtained by doing a permutation on all rows (or all columns) of the identity matrix. The lemma then follows from this. $\square$

Theorem 5.1. Let M_1 and M_2 in Λ_n^{2r} . Suppose that f is an isomorphism between graded rings

$$\mathcal{R}_{M_1} = V_0^{M_1} + \dots + V_{n-2}^{M_1} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \dots)$$

and

$$\mathcal{R}_{M_2} = V_0^{M_2} + \dots + V_{n-2}^{M_2} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \dots).$$

Then there is an element $\sigma \in \mathcal{W}_{2r}$ such that $f = \sum_{i=0}^{\infty} \sigma t^i$ is analytic, where $f(\beta) =$

$$\sum_{i=0}^{\infty} \sigma(v_i) t^i \text{ for } \beta = \sum_{i=0}^{\infty} v_i t^i \in \mathcal{R}_{M_1}.$$

Proof. Since the restriction $f|_{\mathcal{V}_{2r}t^{n-1}} : \mathcal{V}_{2r}t^{n-1} \longrightarrow \mathcal{V}_{2r}t^{n-1}$ is a linear isomorphism, there exists an automorphism σ of $\mathcal{V}_{2r}$ such that $f|_{\mathcal{V}_{2r}t^{n-1}} = \sigma t^{n-1}$.

First, let us show that $\sigma \in \mathcal{W}_{2r}$. Since $x \circ x = x$ for any $x \in (\mathbb{Z}_2)^{2r}$ and f is a ring isomorphism, one has that for any $v \in \mathcal{V}_{2r}$,

$$f(vt^{2n-2}) = f(v \circ vt^{2n-2}) = f(vt^{n-1})f(vt^{n-1}) = \sigma(v) \circ \sigma(v)t^{2n-2} = \sigma(v)t^{2n-2}$$

so $f|_{\mathcal{V}_{2r}t^{2n-2}} = \sigma t^{2n-2}$. Furthermore, for any $v_1, v_2 \in \mathcal{V}_{2r}$, one has that

$$\sigma(v_1 \circ v_2)t^{2n-2} = f(v_1 \circ v_2t^{2n-2}) = f(v_1t^{n-1})f(v_2t^{n-1}) = \sigma(v_1) \circ \sigma(v_2)t^{2n-2}$$

so $\sigma(v_1 \circ v_2) = \sigma(v_1) \circ \sigma(v_2)$. Thus, $\sigma \in \mathcal{W}_{2r}$.

By Theorem 3.1, for each $i < n - 1$, $\underline{1} \in V_i^M \subset \mathcal{V}_{2r}$, and one knows from the proof of Lemma 5.1 that $\sigma(\underline{1}) = \underline{1}$ so $f|_{V_0^M} = \sigma$. By Remark 4 one knows that $\mathcal{R}_{M_i}, i = 1, 2$, are free $\mathbb{Z}_2[t]$ -modules, and it is easy to see that f is also an isomorphism between free $\mathbb{Z}_2[t]$ -modules $\mathcal{R}_{M_1}$ and $\mathcal{R}_{M_2}$. For $0 \leq i < n - 1$, let $v \in V_i^M \subset \mathcal{V}_{2r}$, since $f|_{\mathcal{V}_{2r}t^{n-1}} = \sigma t^{n-1}$, one has that

$$f(vt^i)t^{n-1-i} = f(vt^{n-1}) = \sigma(v)t^{n-1}$$

so $f(vt^i) = \sigma(v)t^i$. Thus, $f|_{V_i^M t^i} = \sigma t^i$ for $0 \leq i < n - 1$.

Let $\ell = a(n - 1) + b \geq n$ with $b < n - 1$. For any $v \in \mathcal{V}_{2r}$, one has that

$$f(vt^\ell) = f(\underbrace{v \circ \cdots \circ v}_a t^{a(n-1)})t^b = [f(vt^{n-1})]^a t^b = \sigma(v) \underbrace{\circ \cdots \circ}_a \sigma(v)t^\ell = \sigma(v)t^\ell.$$

Thus, for $i \geq n$, $f|_{\mathcal{V}_{2r}^M t^i} = \sigma t^i$.

Since $\mathcal{V}_{2r}$ is not closed with respect to the operation $\circ$ by Corollary 3.1, there must be $u, w \in \mathcal{V}_{2r}$ such that $u \circ w \notin \mathcal{V}_{2r}$. Since $\mathcal{V}_{2r}$ has dimension $2r - 1$, one then has that $(\mathbb{Z}_2)^{2r} = \mathcal{V}_{2r} + \text{Span}\{u \circ w\}$. Actually, this is a direct sum decomposition of $(\mathbb{Z}_2)^{2r}$, i.e., $(\mathbb{Z}_2)^{2r} = \mathcal{V}_{2r} \oplus \text{Span}\{u \circ w\}$. Furthermore, let $x \in (\mathbb{Z}_2)^{2r}$, then there is a vector $v \in \mathcal{V}_{2r}$ such that x can be written as $v + \varepsilon u \circ w$ where $\varepsilon = 0$ or 1 . For $i = a(n - 1) + b \geq n$ with $b < n - 1$, since $f|_{\mathcal{V}_{2r}^M t^j} = \sigma t^j$ for any $j \geq n - 1$, one has that

$$\begin{aligned} f(xt^i)t^{n-1-b} &= f(vt^i + \varepsilon u \circ wt^i)t^{n-1-b} \\ &= f(vt^i)t^{n-1-b} + \varepsilon f(u \circ wt^i)t^{n-1-b} \\ &= f(vt^{(a+1)(n-1)}) + \varepsilon f([ut^{n-1}] \circ [wt^{a(n-1)}]) \\ &= \sigma(v)t^{(a+1)(n-1)} + \varepsilon f(ut^{n-1})f(wt^{a(n-1)}) \\ &= \sigma(v)t^{(a+1)(n-1)} + \varepsilon \sigma(u) \circ \sigma(w)t^{(a+1)(n-1)} \\ &= [\sigma(v) + \varepsilon \sigma(u) \circ \sigma(w)]t^{(a+1)(n-1)} \\ &= \sigma(v + \varepsilon u \circ w)t^{(a+1)(n-1)} \text{ since } \sigma \in \mathcal{W}_{2r} \\ &= \sigma(x)t^{i+(n-1-b)} \end{aligned}$$

so $f(xt^i) = \sigma(x)t^i$. Thus $f|_{(\mathbb{Z}_2)^{2r} t^i} = \sigma t^i$ for $i \geq n$.

Combining the above argument, we complete the proof. $\square$

Theorem 5.2. *Let M_1 and M_2 in Λ_n^{2r} . Then $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ are isomorphic if and only if there exists an element $\sigma \in \mathcal{W}_{2r}$ such that σ isomorphically maps $V_i^{M_1}$ onto $V_i^{M_2}$ for $i < n - 1$.*

Proof. Suppose that $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ are isomorphic. Then by Theorem 3.1 there is an isomorphism f between graded rings

$$\mathcal{R}_{M_1} = V_0^{M_1} + \cdots + V_{n-2}^{M_1} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \cdots)$$

and

$$\mathcal{R}_{M_2} = V_0^{M_2} + \cdots + V_{n-2}^{M_2} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \cdots).$$

One knows from Theorem 5.1 that there is an element $\sigma \in \mathcal{W}_{2r}$ such that $f = \sum_{i=0}^{\infty} \sigma t^i$.

Then the restriction $f|_{V_i^{M_1} t^i} = \sigma t^i$, which is an isomorphism from $V_i^{M_1} t^i$ to $V_i^{M_2} t^i$ for $i < n - 1$. Thus σ isomorphically maps $V_i^{M_1}$ onto $V_i^{M_2}$ for $i < n - 1$.

Conversely, if there exists an element $\sigma \in \mathcal{W}_{2r}$ such that σ isomorphically maps $V_i^{M_1}$ onto $V_i^{M_2}$ for $i < n - 1$, then by Theorem 5.1, $\sum_{i=0}^{\infty} \sigma t^i$ gives an isomorphism between graded rings $\mathcal{R}_{M_1} = V_0^{M_1} + \cdots + V_{n-2}^{M_1} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \cdots)$ and $\mathcal{R}_{M_2} = V_0^{M_2} + \cdots + V_{n-2}^{M_2} t^{n-2} + \mathcal{V}_{2r} t^{n-1} + (\mathbb{Z}_2)^{2r} (t^n + \cdots)$. Then $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ are isomorphic by Theorem 3.1. $\square$

Remark 7. Let $M_1, M_2 \in \Lambda_n$. One sees from Theorem 5.2 that if $|M_1^G| \neq |M_2^G|$, then $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ must not be isomorphic.

In the case $n = 3$, for $M \in \Lambda_3^{2r}$, since

$$H_G^*(M; \mathbb{Z}_2) \cong V_0^M + V_1^M t + \mathcal{V}_{2r} t^2 + (\mathbb{Z}_2)^{2r} (t^3 + \cdots),$$

one sees that the structure of $H_G^*(M; \mathbb{Z}_2)$ actually depends upon that of V_1^M . Thus, Theorem 5.2 has a simpler expression in this case.

Corollary 5.1. *Let M_1 and M_2 in Λ_3^{2r} . Then $H_G^*(M_1; \mathbb{Z}_2)$ and $H_G^*(M_2; \mathbb{Z}_2)$ are isomorphic if and only if there exists an element $\sigma \in \mathcal{W}_{2r}$ such that σ isomorphically maps $V_1^{M_1}$ onto $V_1^{M_2}$.*

6. THE NUMBER OF EQUIVARIANT COHOMOLOGY STRUCTURES

In this section we shall consider the number of equivariant cohomology rings up to isomorphism of all 3-dimensional G -manifolds in Λ_3^{2r} . For $M \in \Lambda_3^{2r}$ one knows that V_1^M has dimension r and is the largest-dimensional subspace of $\mathcal{V}_{2r}$ with the property that $u \circ v \in \mathcal{V}_{2r}$ for $u, v \in V_1^M$ by Corollary 3.2.

Let $\mathcal{M}_r$ denote the set of those $2r \times r$ matrices $(v_1, \dots, v_r)$ with rank r over $\mathbb{Z}_2$ such that $v_i \circ v_j \in \mathcal{V}_{2r}$ for any $1 \leq i, j \leq r$. $\mathcal{M}_r$ admits the following two actions.

One action is the right action of $\text{GL}(r, \mathbb{Z}_2)$ on $\mathcal{M}_r$ defined by $(v_1, \dots, v_r) \lambda$ for $\lambda \in \text{GL}(r, \mathbb{Z}_2)$. It is easy to see that such action is free. Obviously, all column vectors of

each matrix $(v_1, \dots, v_r)$ span the same linear space as all column vectors of $(v_1, \dots, v_r)\lambda$ for $\lambda \in \text{GL}(r, \mathbb{Z}_2)$.

The other action is the left action of the Weyl group $\mathcal{W}_{2r} = \mathcal{S}_{2r}$ on $\mathcal{M}_r$ defined by $\sigma(v_1, \dots, v_r) = (\sigma v_1, \dots, \sigma v_r)$ for $(v_1, \dots, v_r) \in \mathcal{M}_r$ and $\sigma \in \mathcal{S}_{2r}$. In general, this action is not free.

For $M \in \Lambda_3^{2r}$, since the space $V_1^M \subset \mathcal{V}_{2r}$ may be spanned by all column vectors of some matrix $(v_1, \dots, v_r)$ in $\mathcal{M}_r$, the number of all possible spaces $V_1^M \subset \mathcal{V}_{2r}$ is at most $|\mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)| = \frac{|\mathcal{M}_r|}{|\text{GL}(r, \mathbb{Z}_2)|}$.

Together with the above understood and Corollary 5.1, one has

Proposition 6.1. *The number of equivariant cohomology rings up to isomorphism of all $\mathbb{Z}_2$ -manifolds in Λ_3^{2r} is at most*

$$|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)|.$$

Remark 8. There is a natural map g from Λ_3^{2r} to $\mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$. If this map is surjective, then the number of equivariant cohomology rings up to isomorphism of all $\mathbb{Z}_2$ -manifolds in Λ_3^{2r} is exactly $|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)|$. To determine whether g is surjective or not is an interesting thing, but it seems to be quite difficult.

Generally, the computation of the number $|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)|$ is not an easy thing. Next, we shall analyze this number.

Taking an element $A \in \mathcal{M}_r$, there must be $\sigma \in \mathcal{S}_{2r}$ and $\lambda \in \text{GL}(r, \mathbb{Z}_2)$ such that

$$\sigma A \lambda = \begin{pmatrix} I_r \\ P \end{pmatrix}$$

so each orbit of the orbit set $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$ contains the representative of the form $\begin{pmatrix} I_r \\ P \end{pmatrix}$, where I_r is the $r \times r$ identity matrix. It is easy to check that $P \in O(r, \mathbb{Z}_2)$, where $O(r, \mathbb{Z}_2)$ is the orthogonal matrix group over $\mathbb{Z}_2$. Obviously, $O(r, \mathbb{Z}_2)$ always admits the left and right actions of the Weyl group $\mathcal{S}_r$ in $\text{GL}(r, \mathbb{Z}_2)$.

Lemma 6.1. $|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)| \leq |\mathcal{S}_r \backslash O(r, \mathbb{Z}_2) / \mathcal{S}_r|$.

Proof. Suppose that $P_1, P_2 \in O(r, \mathbb{Z}_2)$ belong to the same orbit in $\mathcal{S}_r \backslash O(r, \mathbb{Z}_2) / \mathcal{S}_r$. Then there exist τ, ρ in $\mathcal{S}_r$ such that

$$\tau P_1 \rho = P_2.$$

Let $\sigma = \begin{pmatrix} \rho^{-1} & 0 \\ 0 & \tau \end{pmatrix}$. Then $\sigma \in \mathcal{S}_{2r}$. Obviously,

$$\sigma \begin{pmatrix} I_r \\ P_1 \end{pmatrix} \rho = \begin{pmatrix} \rho^{-1} \rho \\ \tau P_1 \rho \end{pmatrix} = \begin{pmatrix} I_r \\ P_2 \end{pmatrix}$$

so $\begin{pmatrix} I_r \\ P_1 \end{pmatrix}$ and $\begin{pmatrix} I_r \\ P_2 \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$. This means that

$$|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)| \leq |\mathcal{S}_r \backslash O(r, \mathbb{Z}_2) / \mathcal{S}_r|.$$

This completes the proof. $\square$

Remark 9. Generally, $|\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)|$ is not equal to $|\mathcal{S}_r \backslash O(r, \mathbb{Z}_2) / \mathcal{S}_r|$. For example, take

$$P_1 = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix} \text{ and } P_2 = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}$$

it is easy to check that P_1 and P_2 don't belong to the same orbit in $\mathcal{S}_6 \backslash O(6, \mathbb{Z}_2) / \mathcal{S}_6$, but $\begin{pmatrix} I_6 \\ P_1 \end{pmatrix}$ and $\begin{pmatrix} I_6 \\ P_2 \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{12} \backslash \mathcal{M}_6 / \text{GL}(6, \mathbb{Z}_2)$.

Definition 6.1. Let $A = (v_1, \dots, v_r) \in \mathcal{M}_r$. One says that A is *irreducible* if the space $\text{Span}\{v_1, \dots, v_r\}$ cannot be decomposed as a direct sum of some nonzero subspaces $V_1, \dots, V_l, l > 1$, with the property that $V_i \circ V_j = \{\underline{0}\}$ for $i \neq j$, where $V_i \circ V_j = \{\underline{0}\}$ means that for any $x \in V_i$ and $y \in V_j$, $x \circ y = \underline{0}$.

We would like to point out that if one can find out all possible irreducible matrices of $\mathcal{M}_r$ for any r , then one can construct a representative of each class in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$, with the form

$$\begin{pmatrix} A_1 & 0 & \dots & 0 \\ 0 & A_2 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & A_s \end{pmatrix}$$

where the blocks A_i 's are irreducible matrices.

The following result shows that there exists at least one irreducible matrix in $\mathcal{M}_r$ for almost any positive integer r . Let

$$A(l) = \begin{pmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & 0 & \dots & 1 & 0 \\ & & & & \dots & & \\ 1 & 0 & 0 & 1 & \dots & 0 & 0 \\ 1 & 0 & 1 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 1 & 1 & \dots & 1 & 1 \\ 0 & 1 & 1 & 1 & \dots & 1 & 1 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 1 & \dots & 0 & 0 \\ & & & & \dots & & \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}_{(2l-1) \times (l-1)}$$

be a $(2l-1) \times (l-1)$ matrix with $l \geq 4$ and $\underline{1}_{l \times 1} = (1, \dots, 1)^\top$. Note that only when $l \geq 4$ is even, the first column vector v of $A(l)$ exactly has the property $|v| = 0$ in $\mathbb{Z}_2$.

Lemma 6.2. (a) For even $r \geq 4$, there exists an irreducible $2r \times r$ matrix

$$\begin{pmatrix} A(r) & \underline{1}_{(2r-1) \times 1} \\ 0 & 1 \end{pmatrix}.$$

(b) For odd $r \geq 7$, there exist irreducible $2r \times r$ matrices of the following form

$$\begin{pmatrix} A(s) & 0 & \underline{1}_{(2s-1) \times 1} \\ 0 & A(t) & \underline{1}_{(2t-1) \times 1} \end{pmatrix}$$

with only two blocks $A(l)$'s, where $s+t = r+1$ and $s, t \geq 4$ are even. In particular, both $\begin{pmatrix} A(s_1) & 0 & \underline{1}_{(2s_1-1) \times 1} \\ 0 & A(t_1) & \underline{1}_{(2t_1-1) \times 1} \end{pmatrix}$ and $\begin{pmatrix} A(s_2) & 0 & \underline{1}_{(2s_2-1) \times 1} \\ 0 & A(t_2) & \underline{1}_{(2t_2-1) \times 1} \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$ if and only if $\{s_1, t_1\} = \{s_2, t_2\}$, where $s_j + t_j = r+1$ and $s_j, t_j \geq 4$ are even for $j = 1, 2$.

Remark 10. (i) A direct observation shows that when $r = 1$, $\mathcal{M}_1$ contains a unique matrix $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$, which is irreducible, and when $r = 2, 3, 5$, there is no any irreducible matrix. However, we don't know whether those irreducible matrices stated in Lemma 6.2, with $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ together, give all possible irreducible matrices.

(ii) For odd $r \geq 7$, let $\lambda(r)$ denote the number of the orbit classes of irreducible matrices in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$. By Lemma 6.2, $\lambda(r)$ is equal to or more than the

number of the solutions (x, y) of the following equation

$$\begin{cases} x + y = \frac{r+1}{2} \\ x \geq y \geq 2. \end{cases}$$

Then, an easy argument shows that the number of solutions is $\lfloor \frac{(r+1)/2-1-1}{2} \rfloor = \lfloor \frac{r-3}{4} \rfloor$, so $\lambda(r) \geq \lfloor \frac{r-3}{4} \rfloor$.

Let $A = (v_1, \dots, v_r) \in \mathcal{M}_r$. Set

$$\mathcal{V}(A) := \text{Span}\{v_1, \dots, v_r\}$$

and

$$\mathcal{X}(A) := \text{Span}\{v_i \circ v_j | i, j = 1, \dots, r\}.$$

It is easy to see the following properties:

- (P1) $\mathcal{V}(A) \subset \mathcal{X}(A) \subset \mathcal{V}_{2r}$ so $r \leq \dim \mathcal{X}(A) \leq 2r - 1$;
- (P2) If $A, B \in \mathcal{M}_r$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$, then $\mathcal{X}(A)$ is linearly isomorphic to $\mathcal{X}(B)$.

Fact 1. *Let $A \in \mathcal{M}_r$. If A is not irreducible, then there exists a $2k \times k$ matrix $A_{2k \times k}$ and a $2l \times l$ matrix $A_{2l \times l}$ with $k + l = r$ such that A and $\begin{pmatrix} A_{2k \times k} & 0 \\ 0 & A_{2l \times l} \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$.*

Proof. If A is not irreducible, by definition, there exists a $k_1 \times k_2$ matrix $A_{k_1 \times k_2}$ and a $l_1 \times l_2$ matrix $A_{l_1 \times l_2}$ with $k_1 + l_1 = 2r$ and $k_2 + l_2 = r$ such that A and $\begin{pmatrix} A_{k_1 \times k_2} & 0 \\ 0 & A_{l_1 \times l_2} \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$. By Proposition 3.1 and the proof method of Corollary 3.1, one has that $k_2 \leq k_1/2$ and $l_2 \leq l_1/2$. Then the relations $k_1 + l_1 = 2r$ and $k_2 + l_2 = r$ force $k_2 = k_1/2$ and $l_2 = l_1/2$. $\square$

Fact 2. *Let $A \in \mathcal{M}_r$. If $\dim \mathcal{X}(A) = 2r - 1$, then A is irreducible.*

Proof. Suppose that A is not irreducible. By Fact 1, there is a matrix $\begin{pmatrix} A_{2k \times k} & 0 \\ 0 & A_{2l \times l} \end{pmatrix}$ that is in the same orbit as A . Thus

$$\dim \mathcal{X}(A) = \dim \mathcal{X}(A_{2k \times k}) + \dim \mathcal{X}(A_{2l \times l}) \leq (2k - 1) + (2l - 1) = 2r - 2$$

by (P1). This is a contradiction. $\square$

Remark 11. Let $A \in \mathcal{M}_r$. Then there is a matrix $P \in O(r, \mathbb{Z}_2)$ such that A and $\begin{pmatrix} I_r \\ P \end{pmatrix}$ belong to the same orbit in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$. Furthermore, an easy argument shows by (P2) that if $\dim \mathcal{X}(A) = r$, then P can be chosen as being I_r , so A is not irreducible when $r \neq 1$.

Now let us give the proof of Lemma 6.2.

Proof of Lemma 6.2. Let $A = \begin{pmatrix} A(r) & \mathbb{1}_{(2r-1) \times 1} \\ 0 & 1 \end{pmatrix} = (v_1, \dots, v_{r-1}, \mathbb{1}_{2r \times 1})$. By Fact 2, it suffices to check that $\dim \mathcal{X}(A) = 2r - 1$. A direct observation shows that

$$\{v_1 \circ v_2, v_1 \circ v_3, \dots, v_1 \circ v_{r-1}, v_2 \circ v_3, v_1, v_2, v_3, \dots, v_{r-1}, \mathbb{1}_{2r \times 1}\}$$

are linearly independent, so $\dim \mathcal{X}(A) \geq 2r - 1$. Since $\dim \mathcal{X}(A) \leq 2r - 1$ by (P1), $\dim \mathcal{X}(A) = 2r - 1$. This completes the proof of Lemma 6.2(a).

In a similar way to the above argument, one may show that $\begin{pmatrix} A(s) & 0 & \mathbb{1}_{(2s-1) \times 1} \\ 0 & A(t) & \mathbb{1}_{(2t-1) \times 1} \end{pmatrix}$ is irreducible, too.

Suppose that $\begin{pmatrix} A(s_1) & 0 & 0 & \dots & 0 & \mathbb{1}_{(2s_1-1) \times 1} \\ 0 & A(s_2) & 0 & \dots & 0 & \mathbb{1}_{(2s_2-1) \times 1} \\ 0 & 0 & A(s_3) & \dots & 0 & \mathbb{1}_{(2s_3-1) \times 1} \\ & & & \dots & & \\ 0 & 0 & 0 & \dots & A(s_k) & \mathbb{1}_{(2s_k-1) \times 1} \end{pmatrix}$ is a $2r \times r$ matrix and

is irreducible. Then one must have that $\sum_{i=1}^k (s_i - 1) + 1 = r$ and $\sum_{i=1}^k (2s_i - 1) = 2r$. From these two equations, it is easy to check that k must be 2.

The proof of the last part of Lemma 6.2(b) is immediate. $\square$

Remark 12. By a direct computation, one may give the first 6 numbers for $N(r) = |\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)|$ as follows:

r	1	2	3	4	5	6
$N(r)$	1	1	1	2	2	3

The representatives of all orbits in $\mathcal{S}_{2r} \backslash \mathcal{M}_r / \text{GL}(r, \mathbb{Z}_2)$ for $r \leq 6$ are stated as follows:

r	1	2	3	4	5	6
Rep.	$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$\begin{pmatrix} I_2 \\ I_2 \end{pmatrix}$	$\begin{pmatrix} I_3 \\ I_3 \end{pmatrix}$	$\begin{pmatrix} I_4 \\ I_4 \end{pmatrix}, B(4)$	$\begin{pmatrix} I_5 \\ I_5 \end{pmatrix}, \begin{pmatrix} B(4) & 0 \\ 0 & \mathbb{1}_{2 \times 1} \end{pmatrix}$	$\begin{pmatrix} I_6 \\ I_6 \end{pmatrix}, \begin{pmatrix} B(4) & 0 & 0 \\ 0 & \mathbb{1}_{2 \times 1} & 0 \\ 0 & 0 & \mathbb{1}_{2 \times 1} \end{pmatrix}, B(6)$

where $B(l) = \begin{pmatrix} A(l) & \mathbb{1}_{(2l-1) \times 1} \\ 0 & 1 \end{pmatrix}$. Note that for $r \geq 7$, one also can find the lower bound of $N(r)$ as follows:

r	7	8	9	10	11	12	13	14	$\dots$
$N(r) \geq$	4	6	7	9	12	16	20	25	$\dots$

REFERENCES

- [AB1] J.L. Alperin and R.B. Bell, *Groups and representations*, Graduate Texts in Mathematics **162** (1995), Springer-Verlag.

- [AB2] M. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology **23** (1984), 1-28.
- [AP] C. Allday and V. Puppe, *Cohomological Methods in Transformation Groups*, Cambridge Studies in Advanced Mathematics, **32**, Cambridge University Press, 1993.
- [B] G. E. Bredon, *Introduction to compact transformation groups*, Pure and Applied Mathematics, Vol. **46** Academic Press, New York-London, 1972.
- [BGH] D. Biss, V. Guillemin and T. S. Holm, *The mod 2 cohomology of fixed point sets of anti-symplectic involutions*, Adv. Math. **185** (2004), 370-399.
- [BV] N. Berline and M. Vergne, *Classes caractéristiques équivariantes. Formules de localisation en cohomologie équivariante*, C. R. Acad. Sci. Paris Sr. I Math. **295** (1982), 539-541.
- [CF] P.E. Conner and E.E. Floyd, *Differentiable periodic maps*, Ergebnisse Math. Grenzgebiete, N. F., Bd. **33**, Springer-Verlag, Berlin, 1964.
- [CS] T. Chang and T. Skjelbred, *The topological Schur lemma and related results*, Ann. of Math. **100** (1974), 307-321.
- [D] T. tom Dieck, *Characteristic numbers of G -manifolds. I*, Invent. Math. **13** (1971), 213-224.
- [GH] R. Goldin and T. S. Holm, *The equivariant cohomology of Hamiltonian G -spaces from residual S^1 actions*, Math. Res. Lett. **8** (2001), 67-77.
- [GKM] M. Goresky, R. Kottwitz, and R. MacPherson *Equivariant cohomology, Koszul duality, and the localization theorem*, Invent. Math. **131** (1998), 25-83.
- [GZ] V. Guillemin and C. Zara, *1-Skeleta, Betti numbers, and equivariant cohomology*, Duke Math. J. **107** (2001), 283-349.
- [H] W. Y. Hsiang, *Cohomology theory of topological transformation groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band **85**. Springer-Verlag, New York-Heidelberg, 1975.
- [KS] C. Kosniowski and R.E. Stong, *$(\mathbb{Z}_2)^k$ -actions and characteristic numbers*, Indiana Univ. Math. J. **28** (1979), 723-743.
- [KP] M. Kreck and V. Puppe, *Involutions on 3-manifolds and self-dual, binary codes*, arXiv:0707.1599.
- [L] Z. Lü, *Graphs and $(\mathbb{Z}_2)^k$ -actions*, arXiv: math.AT/0508643.
- [P] V. Puppe, *Group actions and codes*, Canad. J. math. **153** (2001), 212-224.

SCHOOL OF MATHEMATICAL SCIENCE, FUDAN UNIVERSITY, SHANGHAI, 200433, PEOPLE'S REPUBLIC OF CHINA.

E-mail address: 061018005@fudan.edu.cn

INSTITUTE OF MATHEMATICS, SCHOOL OF MATHEMATICAL SCIENCE, FUDAN UNIVERSITY, SHANGHAI, 200433, PEOPLE'S REPUBLIC OF CHINA.

E-mail address: zlu@fudan.edu.cn